

ISA Atmospheric Model

Jorge Martinez

Compute temperature, pressure, density, and speed of sound from sea level to 86 km using the International Standard Atmosphere (ISA, ISO 2533:1975). Apply the model layer by layer, then plot each property against altitude.

```
import numpy as np
import matplotlib.pyplot as plt
```

ISA Layer Definitions

Define each layer by its base geopotential altitude in meters, base temperature in kelvin, and temperature lapse rate in kelvin per meter.

```
LAYERS = [
    # (base altitude m, base temperature K, lapse rate K/m)
    (0, 288.15, -0.0065), # Troposphere
    (11_000, 216.65, 0.0), # Tropopause (isothermal)
    (20_000, 216.65, +0.001), # Lower stratosphere
    (32_000, 228.65, +0.0028), # Upper stratosphere
    (47_000, 270.65, 0.0), # Stratopause (isothermal)
    (51_000, 270.65, -0.0028), # Lower mesosphere
    (71_000, 214.65, -0.002), # Upper mesosphere
    (86_000, 186.87, 0.0), # Mesopause boundary (limit)
]

# Physical constants
R = 287.058 # Specific gas constant for dry air, J/(kg·K)
g0 = 9.80665 # Standard gravitational acceleration, m/s2
gamma = 1.4 # Ratio of specific heats for dry air

# Sea-level reference values
T0 = 288.15 # K
P0 = 101_325 # Pa
rho0 = P0 / (R * T0)
```

Compute ISA Properties

Start with sea-level pressure. For each altitude, find the layer and compute the temperature from its base altitude and lapse rate:

$$T(h) = T_b + L(h - h_b)$$

The subscript b denotes the base of the current layer. In an isothermal layer, use the exponential pressure relation:

$$P(h) = P_b \exp\left(-\frac{g_0(h - h_b)}{RT_b}\right)$$

For a layer with a nonzero lapse rate, use:

$$P(h) = P_b \left(\frac{T(h)}{T_b}\right)^{-g_0/(RL)}$$

Then compute density and speed of sound:

$$\rho(h) = \frac{P(h)}{RT(h)}$$

$$a(h) = \sqrt{\gamma RT(h)}$$

R is the specific gas constant for dry air, g0 is standard gravitational acceleration, and gamma is the ratio of specific heats.

```
def isa_properties(altitudes_m: np.ndarray):  
    """  
        Compute ISA temperature, pressure, density, and speed of sound.  
  
        Parameters  
        -----  
        altitudes_m : numpy.ndarray  
            Geopotential altitudes in meters.  
  
        Returns  
        -----  
        T : numpy.ndarray  
            Temperature in K.
```

```

P : numpy.ndarray
    Pressure in Pa.
rho : numpy.ndarray
    Air density in kg/m3.
a : numpy.ndarray
    Speed of sound in m/s.

"""
T = np.empty_like(altitudes_m)
P = np.empty_like(altitudes_m)

for i, h in enumerate(altitudes_m):
    # Find the layer containing altitude h.
    P_base = P0
    for layer_idx in range(len(LAYERS) - 1):
        h_base, T_b, L = LAYERS[layer_idx]
        h_top = LAYERS[layer_idx + 1][0]

        if h <= h_top:
            dh = h - h_base
            T[i] = T_b + L * dh
            if L == 0.0:
                P[i] = P_base * np.exp(-g0 * dh / (R * T_b))
            else:
                P[i] = P_base * (T[i] / T_b) ** (-g0 / (L * R))
            break
        else:
            # Compute pressure at the next layer boundary.
            dh = h_top - h_base
            T_top = T_b + L * dh
            if L == 0.0:
                P_base = P_base * np.exp(-g0 * dh / (R * T_b))
            else:
                P_base = P_base * (T_top / T_b) ** (-g0 / (L * R))

    rho = P / (R * T)
    a = np.sqrt(gamma * R * T)
    return T, P, rho, a

# Altitude grid from sea level to 86 km
h = np.linspace(0, 86_000, 1_000)

T, P, rho, a = isa_properties(h)

```

```
h_km = h / 1_000 # Convert to km for plotting.
```

Plot Atmospheric Properties


Plot each property against altitude. Shade the layers to show where the lapse rate changes.

```
fig, axes = plt.subplots(1, 4, figsize=(14, 6), sharey=True)
fig.suptitle("International Standard Atmosphere (ISA)", fontsize=14, fontweight="bold")
# Shade the ISA layers.
layer_colors = [
    "#e8f4f8",
    "#d0eaf4",
    "#b8dff0",
    "#9fd4ec",
    "#87c9e8",
    "#6fbee4",
    "#57b3e0",
    "#3fa8dc",
]
for ax in axes:
    for k in range(len(LAYERS) - 1):
        h_bot = LAYERS[k][0] / 1_000
        h_top = LAYERS[k + 1][0] / 1_000
        ax.axhspan(
            h_bot,
            h_top,
            color=layer_colors[k % len(layer_colors)],
            alpha=0.25,
            zorder=0,
        )
axes[0].plot(T, h_km, color="tab:red", linewidth=1.8)
axes[0].set_xlabel("Temperature (K)")
axes[0].set_ylabel("Altitude (km)")
axes[0].set_title("Temperature")
axes[0].grid(True, linestyle="--", alpha=0.5)
axes[1].plot(P / 1_000, h_km, color="tab:blue", linewidth=1.8)
axes[1].set_xlabel("Pressure (kPa)")
axes[1].set_title("Pressure")
axes[1].grid(True, linestyle="--", alpha=0.5)
axes[2].plot(rho, h_km, color="tab:green", linewidth=1.8)
axes[2].set_xlabel("Density (kg/m³)")
axes[2].set_title("Density")
```

```

axes[2].grid(True, linestyle="--", alpha=0.5)
axes[3].plot(a, h_km, color="tab:orange", linewidth=1.8)
axes[3].set_xlabel("Speed of sound (m/s)")
axes[3].set_title("Speed of sound")
axes[3].grid(True, linestyle="--", alpha=0.5)
for ax in axes:
    ax.set_ylim(0, 86)
plt.tight_layout()
plt.show()

```



Sea-Level Reference Values

Evaluate the model at sea level and print the reference values with units.

```

T_sl, P_sl, rho_sl, a_sl = [x[0] for x in isa_properties(np.array([0.0]))]
print(f"Sea-level temperature : {T_sl:.2f} K ({T_sl - 273.15:.2f} °C)")
print(f"Sea-level pressure : {P_sl:.2f} Pa ({P_sl / 1e3:.4f} kPa)")
print(f"Sea-level density : {rho_sl:.4f} kg/m³")
print(f"Sea-level speed of sound: {a_sl:.2f} m/s")

```

```

Sea-level temperature : 288.15 K (15.00 °C)
Sea-level pressure : 101325.00 Pa (101.3250 kPa)
Sea-level density : 1.2250 kg/m³
Sea-level speed of sound: 340.30 m/s

```