

Monte Carlo Estimation of

Jorge Martinez

Estimate π by sampling points uniformly from a square that contains a unit circle. The square has side length 2, so the ratio of their areas is:

$$\frac{A_{\text{circle}}}{A_{\text{square}}} = \frac{\pi}{4}$$

Use the fraction of points inside the circle to estimate this ratio.

```
import numpy as np
import matplotlib.pyplot as plt
```

Simulation Setup

Set a fixed random seed so repeated runs use the same samples.

```
rng = np.random.default_rng(seed=42)
N = 50_000 # total number of random points
```

Run the Simulation

Draw 50,000 points uniformly from the square:

$$(x, y) \in [-1, 1] \times [-1, 1]$$

A point lies inside the unit circle when:

$$x^2 + y^2 \leq 1$$

Count those points and compute the estimate:

$$\hat{\pi} = 4 \frac{N_{\text{inside}}}{N}$$

```
x = rng.uniform(-1, 1, N)
y = rng.uniform(-1, 1, N)
inside = x**2 + y**2 <= 1.0

pi_estimate = 4 * inside.sum() / N
print(
    f" estimate after {N:,} samples: {pi_estimate:.6f} (error: {abs(pi_estimate - np.pi):.2f})
")
```

```
estimate after 50,000 samples: 3.146320 (error: 4.73e-03)
```

Convergence of the Estimate

Compute a running estimate after each sample. The count of points inside the circle accumulates as more points are drawn:

$$\hat{\pi}_n = 4 \frac{N_{\text{inside}}(n)}{n}$$

```
sample_counts = np.arange(1, N + 1)
running_pi = 4 * np.cumsum(inside) / sample_counts
```

Plot Samples and Convergence

Show the first 5,000 points on the left, colored by whether they fall inside the circle. Plot the running estimate on the right and compare it with π .


```
fig, (ax_scatter, ax_conv) = plt.subplots(1, 2, figsize=(12, 5))
fig.suptitle("Monte Carlo estimation of ", fontsize=14, fontweight="bold")
# Use the first 5,000 points to keep the scatter plot readable.
n_scatter = 5_000
ax_scatter.scatter(
    x[:n_scatter][inside[:n_scatter]],
    y[:n_scatter][inside[:n_scatter]],
    s=0.8,
    color="tab:blue",
    alpha=0.5,
    label="Inside circle",
)
```

```

ax_scatter.scatter(
    x[:n_scatter][~inside[:n_scatter]],
    y[:n_scatter][~inside[:n_scatter]],
    s=0.8,
    color="tab:red",
    alpha=0.5,
    label="Outside circle",
)
theta = np.linspace(0, 2 * np.pi, 300)
ax_scatter.plot(np.cos(theta), np.sin(theta), color="black", linewidth=1.2)
ax_scatter.set_aspect("equal")
ax_scatter.set_title(f"Random samples (n = {n_scatter:,})")
ax_scatter.set_xlabel("x")
ax_scatter.set_ylabel("y")
ax_scatter.legend(loc="lower right", markerscale=6, fontsize=8)
ax_scatter.grid(True, linestyle="--", alpha=0.4)
# Plot the running estimate against the number of samples.
ax_conv.semilogx(
    sample_counts, running_pi, color="tab:blue", linewidth=1.2, label="Running estimate"
)
ax_conv.axhline(np.pi, color="black", linewidth=1.0, linestyle="--", label="True ")
ax_conv.set_title("Convergence of estimate")
ax_conv.set_xlabel("Number of samples")
ax_conv.set_ylabel(" estimate")
ax_conv.legend(fontsize=9)
ax_conv.grid(True, linestyle="--", alpha=0.4)
plt.tight_layout()
plt.show()

```

Monte Carlo estimation of π



Estimate Error

Compare the final estimate with NumPy's value of `np.pi`.

```
print(f"True                : {np.pi:.6f}")
print(f"Estimated          : {pi_estimate:.6f}")
print(f"Absolute error     : {abs(pi_estimate - np.pi):.2e}")
print(f"Relative error     : {abs(pi_estimate - np.pi) / np.pi * 100:.4f} %")
```

```
True                : 3.141593
Estimated          : 3.146320
Absolute error     : 4.73e-03
Relative error     : 0.1505 %
```